9500085

THE UNITED STATES OF ANTERIOA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Western Plant Breeders, Inc.

MICTORS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED, PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE GHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY; OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR ORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT ED BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

BARLEY

'Merlin'

In Testimonn Murreot, I have hereunto set my hand and caused the seal of the Hant Bariety Hrotertian Office to be affixed at the City of Washington, D.C. this thirty-first day of August in the year of our Lord one thousand nine hundred and ninety-five.

Allest

RAARQ

Acting Commissioner Plant Variety Protection Of

Plant Variety Protection Office Agricultural Marketing Service

Secretary of Agriculture

| TIEF HODOGE EGGNEETT MOIDE TOTAL TOTAL ETTE E | | op: + = = = = = = = = = = = = = = = = = = | | |
|---|--|---|--|--|
| U.S. DEPARTMENT (AGRICULTURAL MAI SCIENCE I | | Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. | | |
| APPLICATION FOR PLANT VAR | 2421). Information is held contidential until certificate is issued (7 U.S.C. 2426). | | | |
| NAME OF APPLICANT(S) (as it is to appear on the Certificate | | TEMPORARY DESIGNATION OR EXPERIMENTAL NO. | 3. VARIETY NAME | |
| Western Plant Breeders, Inc. | | BZ 489-29 | Merlin | |
| 4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP) | | 5. PHONE (include area code) | FOR OFFICIAL USE ONLY | |
| 8111 Timberline Drive | | (406) 587-1218 | 9500035 | |
| Bozeman, Montana 59715 | | | Date Mar. 30, 1995 | |
| 6. GENUS AND SPECIES NAME | 7. FAMILY NAME (Bot | tanical | Time N □ A.M. □ P.M. | |
| Hordeum vulgare | Graminea | · | G 1 | |
| | GrainThea | 9. DATE OF DETERMINATION | E S 0 225 00 | |
| 8. CROP KIND NAME (Common Name) Barley | | | S Date | |
| · · · · · · · · · · · · · · · · · · · | LOE ODGANIZATION | May 1, 1990 | R Mar. 30, 1995 | |
| 10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FOR association, etc.) | M OF ORGANIZATION | (Corporation, partnership, | C Certificate Fee: | |
| Corporation | | | \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\ | |
| 11. IF INCORPORATED, GIVE STATE OF INCORPORATION | | 12. DATE OF INCORPORATION | D Date | |
| Arizona 13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), | | May 24, 1990 | July 19, 1995 | |
| Western Plant Breede 8111 Timberline Driv Bozeman, MT 59715 14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUB! a Exhibit A, Origin and Breeding History of the Variet b Exhibit B, Novelty Statement c Exhibit C, Objective Description of Variety d Exhibit D, Additional Description of Variety e Exhibit E, Statement of the Basis of Applicant's Ow f. Seed Sample (2.500 viable untreated seeds). Date g. Filing and Examination Fee (\$2,325) made payable | E ATTED (Follow INSTRU y mership Seed Sample mailed to | PHONE (Include area code): UCTIONS on reverse) o Plant Variety Protection OfficeMa | 6) 587-1218 rch 29, 1995 | |
| 15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VA Plant Variety Protection Act) | | | · · · · · · · · · · · · · · · · · · · | |
| 16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS? | 17. IF "YI | | PRODUCTION BEYOND BREEDER SEED? | |
| ☐ YES ☐ NO 18. DID THE APPLICANT(S) PREVIOUSLY FILE FOR PROTECTION | N OF THE VARIETY IN | FOUNDATION REGISTE | RED CERTIFIED | |
| ☐ YES (If "YES," through ☐ Plant Variety Protecti | | It Act. Give date: |). | |
| 19. HAS THE VARIETY BEEN RELEASED, USED, OFFERED FOR | | IN THE U.S. OR OTHER COUNTRIES? | | |
| YES (If "YES," GIVE NAMES OF COUNTRIES AND DA | ies) <u>Hartin</u> | on 100.33 | | |
| The applicant(s) declare(s) that a viable sample of basic seed such regulations as may be applicable. | s of this variety will be | furnished with the application and will be | replenished upon request in accordance with | |
| The undersigned applicant(s) is (are) the owner(s) of this sexus in section 41, and is entitled to protection under the provision | | | y is distinct, uniform, and stable as required | |
| Applicant(s) is (are) informed that false representation herein | ೯೯೮- can jeopardize protectio | on and result in penalties. | | |
| SIGNATURE OF APPLICANT [Owner(s)] | | CAPACITY OR TITLE | DATE | |
| Oale R. Plank | F | Barley&Wheat Breed | er March 29, 1995 | |
| SIGNATURE OF APPLICANT [Owner(s)] | | CAPACITY OR TITLE | DATE | |
| | 1 | | | |
| | • | | | |

14a. Origin and Breeding History

MERLIN is a two-rowed, hulless, waxy, spring barley that was selected by Western Plant Breeders (WPB) from a male-sterile facilitated, recurrent selection population (MSFRSP) for short strawed, two-rowed, waxy, hulless barley. The short, two-rowed population was developed by crossing WPB's two-rowed MSFRSP with a short two-rowed line PH 582-8 in 1982. (PH 582-8 was a selection from the cross of WPB short six-rowed population by various disease resistant lines, including two-rowed barley's.) Short, two-rowed, male sterile F2 plants from the 1982 cross were intercrossed in 1983. Several short, F2 male sterile plants from this intercross were crossed with two lines, EK-8 (an Erbet x Klages selection) and TR-451 (Klages x S72204 selection) in 1984. Short, F2 male sterile plants in this population were intercrossed in 1985. Short F2 male-sterile plants from this intercross were crossed with WPB's two-rowed, waxy hulless line BZ 585-85 (WestBred Waxbar x TR-451) in 1986. Spikes were selected from short, hulless, two-rowed F2 plants in 1987 at Phoenix, AZ and threshed in bulk. The F3 seed was thin planted at Blackfoot, Idaho in 1988 and single F3 plants were selected at harvest. Seed of these plants were planted near Phoenix in the fall of 1988 in 5'x 20' plots. Five F4 plants were selected from each desirable plot in May of 1989 and F5 seed from these were planted in 5' x 20' plots near Bozeman, MT in May, 1989. One such plot was selected, bulk harvested, and given the experimental number, BZ 489-29 in September of 1989. Successive F6, F7, F8, F9 and F10 generation seed from this plot were yield tested in 1990 through 1994 in Montana, Idaho, and Washington.

F6 seed was planted near Phoenix in the fall of 1989 for purification. Fifteen F6 plants were selected in May of 1990 and F7 seed from these plants were planted in individual 5' x 20' plots near Bozeman in May of 1990. Uniform plots were bulk harvested. F8 seed from these plots were planted in 1991 near Bozeman on

approximately one eighth of an acre. F9 seed from this increase were harvested and designated "Breeders" Seed. This F9 seed was again increased as Breeders seed near Bozeman in 1992 and given the name "Merlin". This Breeders seed was planted in May, 1993 and the subsequent production was harvested in September, 1993 as Foundation seed. The Foundation seed was planted in May, 1994 and harvested as Registered and Certified seed in September, 1994. The first sale of Certified seed to the general public will be in the spring of 1995.

Merlin is a stable and uniform variety in agronomic appearance and performance across several generations and growing conditions. No variants are known to occur at this time. Agronomic data to support this stability are presented in Tables 1 through 7.

14b. Novelty Statement

Merlin is most similar to WestBred Waxbar. However, Merlin has long awns while WestBred Waxbar has short awns. Merlin is a semidwarf and WestBred Waxbar is a standard height variety (Merlin is approximately 19 cm shorter than WestBred Waxbar). Also, the spike of Merlin is coated with a wax coating while the spike of WestBred Waxbar is glossy.

A20 V

The above comparisons, along with the complete objective description (14c.), show Merlin to be a novel variety of spring barley.

FORM GR-470-5 (11-1-72)

UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE GRAIN DIVISION

EXHIBIT C (Barley)

HYATTSVILLE, MARYLAND 20782

OBJECTIVE DESCRIPTION OF VARIETY

| INSTRUCTIONS: See Reverse. BARLEY (HORDEUM VULGARE) | |
|---|---|
| Western Plant Breeders, Inc. | FOR OFFICIAL USE ONLY |
| ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code) 8111 Timberline Drive | 9500085 |
| Bozeman, MT 59715 | DESIGNATION Merlin |
| Place the appropriate number that describes the varietal character of this variety in the Place a zero in first box (i.e. 089 or 09) when number is either 99 or less | |
| 1. GROWTH HABIT: | or y or less. |
| | : 1 = PROSTRATE 2 = SEMIPROSTRATE 3 = ERECT |
| 2. MATURITY (50% Flowering): | |
| 2 1 = EARLY (California Mariout) 2 = MIDSEASON (Betzes) 3 = LATE (Frontier) | |
| No. of days Earlier than 8 1 = BETZES 2 = CALIFORNIA MARIOUT | 3 = CONQUEST 4 = DICKSON |
| | w 8= klages |
| 3, PLANT HEIGHT (From soil level to top of head): | |
| 1 1 = SEM!DWARF 2 = SHORT (California Mariout) 3 = MEDIUM TALL (Betzes) | 4 = TALL (Conquest) |
| 2 2 Cm. Shorter than 5 1 = BETZES 2 = CALIFORNIA MARIOUT | 3 = CONQUEST 4 = DICKSON |
| 0 0 Cm. Taller than x 5 = PIROLINE 6 = PRIMUS 7 = UNITA | N x= none listed |
| 4. STEM: | |
| 1 = 0 - 3 cm. 2 = 3 - 10 cm. 1 = 0 - 3 cm. 2 = 3 - 10 cm. 1 = 0 - 3 cm. 2 = 3 - 10 cm. Anthocyanin: | 1 = ABSENT 2 = PRESENT |
| 0 4 NO. OF NODES (Originating from node above ground) Skin i | s waxy appearence |
| 1 Collar Shape: 1 = CLOSED 2 = V-SHAPED 3 = OPEN 1 Shape of Neck: | 1 = STRAIGHT 2 = SNAKY 3 = OTHER (Specify) |
| 5. LEAF: 1 Basal leaf sheath (seedling): 1 = GLABROUS 2 = PUBESCENT 1 Position of flag le | 1 = DROOPING ext(at boot stage): 2 = UPRIGHT |
| 2 Waxiness: 1 = ABSENT (Glossy) 2 = SLIGHTLY WAXY 1 5 MM. WIDTH | (First leaf below flag leaf) |
| 2 3 CM. LENGTH (First leaf below fleg leaf) 1 Anthocyanin in le | af sheath: 1 = ABSENT 2 = PRESENT |
| 6. HEAD: | |
| Type: 1 = TWO BOWED 2 - SIX DOWED | LAX 2 = ERECT (Not dense) ERECT (Dense) |
| | ABSENT (Glossy) 2 = SLIGHTLY WAXY |
| 1 = NONE 2 = AT TIP 3 = 1/4 - 1/2 OF HEAD Rachis (Hair on ed | ge): 1 = LACKING 2 = FEW 3 = COVERED |
| 7. GLUME: | |
| 2 Length: 3 = MORE THAN 1/2 OF LEMMA 1 Hairs: 1 = NON | E 2 = SHORT 3 = LONG |
| Hair covering: 1 = NONE 2 = RESTRICTED TO MIDDLE 3 = CONFINED TO BA | ND 4 - COMPLETELY COVERED |
| Awns: 1 = LESS THAN EQUAL TO LENGTH OF GLUMES 2 = EQUAL TO LENGTH 3 = MORE THAN EQUAL TO LENGTH OF GLUMES | OF GLUMES |
| 2 Awn Surface: 1 = SMOOTH 2 = SEMISMOOTH 3 = ROUGH | |
| | |

| 8. LEMMA: | |
|--|---|
| | OWS AWNLESS ON LATERAL ROWS ATERAL ROWS 4 = SHORT (less than equal to length of spike) |
| 1 Awn Surface: 0 = AWNLESS 1 = SMOOTH 2 = SEMISM | моотн з=Rough rough at the tip |
| Teeth: 1 = ABSENT 2 = FEW 3 = NUMEROUS | 1 Hair: 1 = ABSENT 2 = PRESENT |
| 1 = DEPRESSION 2 = SLIGHT CREASE 3 = TRANSVERSE CREASE | 1 Rachilla Hairs: 1 = SHORT 2 = LONG |
| 9. STIGMA: | |
| 1 Hairs: 1 = FEW 2 = MANY | |
| 10. SEED: | |
| Type: 1 = NAKED 2 = COVERED | Hairs on Ventral Furrow: 1 = ABSENT 2 = PRESENT |
| Length: 1 = SHORT (8.0 mm.) 2 = SHORT TO MIDLONG 4 = MIDLONG TO LONG (9.0 - 10.5 mm.) | (7.5 - 9.0 mm.) 3 = MIDLONG (8.5 - 9.5 mm.) 5 = LONG (10.0 mm.) |
| 1 Wrinkling of hull: 1 = NAKED 2 = SLIGHTLY WRINKLES | 3 = SEMIWRINKLED 4 = WRINKLED |
| Aleurone Color: 1 = COLORLESS (White or Yellow) 2 = B | LUE |
| 0 2 PERCENT ABORTIVE | 4 5 GMS, PER 1000 SEEDS |
| 11. DISEASE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) | |
| 0 SEPTORIA 0 NET BLOTCH | 0 SPOT BLOTCH 1 POWDERY MILDEW |
| 0 LOOSE SMUT 0 BACTERIAL BLIGHT | 0 COVERED SMUT 0 FALSE LOOSE SMUT |
| 1 STEM RUST 0 LEAF RUST | 0 SCAB 1 SCALD |
| 0 AY 0 BSMV | 0 BYDV 1 OTHER (Specify) Strip rust |
| 12. INSECT: (0 = Not tested, 1 = Susceptible 2 = Resistant) | |
| 0 GREEN BUG 0 ENGLISH GRAIN APHID | 0 CHINCH BUG 0 ARMYWORM |
| 0 GRASS HOPPERS 0 CERIAL LEAF BETTLE | O OTHER (Specify) |
| HESSIAN FLY RACES | 0 _B 0 _C |
|) O D O E | 0 F 0 G |
| 3. CHEMICAL (0 = Not Tested, 1 = Susceptible, 2 = Resistant) | |
| 0 DDT 0 OTHER (Specify) | |
| 4. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THAT | SUBMITTED: |
| CHARACTER NAME OF VARIETY | CHARACTER NAME OF VARIETY |
| Plant tillering of WestBred Waxbar | Seed size WestBred Waxbar |
| Leaf size WestBred Waxbar | Coleoptile elongation WestBred Waxbar |
| Leaf color WestBred Waxbar | Seedling pigmentation WestBred Waxbar |
| Leaf carriage WestBred Waxbar | WestBred Waxbar |
| | HODODICA MAXVAI |

REFERENCES: The following publications may be used as a reference aid for the standardization of character descriptions and terms used in this form:

- 1. Wiebe, G. A., and D. A. Reid, 1961, Classification of Barley Varieties Grown in the United States and Canada in 1958, Technical Bulletin No. 1224, U.S. Dept. of Agriculture.
- Reid, D. A., and G. A. Wiebe, 1968, Barley: Origin, Botany, Culture, Winter Hardiness, Genetics, Utilization, Pests, Agriculture Handbook No. 338, U.S. Dept. of Agriculture. pp. 61-84.
- 3. Malting Barley Improvement Association, Milwaukee, Wisconsin, 1971, Barley Variety Dictionary.

COLOR: Nickerson's or any recognized color fan may be used to determine color of the described variety.

9500085

MERLIN

14c. Objective Description (see pages 4 and 5)

14d. Additional Description

Merlin , like WestBred Waxbar, is considered a "low amylose" or "waxy" endosperm variety. The starch composition of the endosperm is mostly amylopectin starch with very little (< 5%) amylose starch. Therefore, the endosperm does not stain blue when Iodine Potassium Iodide (I₂KI) is added.

14e. Statement of Ownership

'Merlin', the variety for which Plant Variety Protection is hereby sought, was developed by Dr. Dale R. Clark, an employee of Western Plant Breeders, Inc.. All rights to any invention, discovery, or development made by the employee while employed by Western Plant Breeders, Inc. were assigned by Western Plant Breeders, Inc. with no rights of any kind pertaining to 'Merlin' are retained by the employees.

Table 1. Heading date (from Jan. 1) of Merlin compared to WestBred Waxbar in Western Plant Breeders, Inc. trials at Bozeman, MT from 1990 to 1994.

| | <u>Merlin</u> | WestBred <u>Waxbar</u> |
|------|---------------|---------------------------|
| 1990 | 186 | 190 |
| 1991 | 192 | 193 |
| 1992 | 174 | 182 |
| 1994 | <u>182</u> | <u>184</u> |
| mean | 183.5 | 187.25 |

900

Table 2. Agronomic data comparing Merlin to WestBred Waxbar in Western Plant Breeders, Inc trials from 1990 to 1994.

| | · - | | Merlin | | <u> </u> | WestBred Waxbar | |
|-------|--------------------|--------------------|--------------------|------------------------|---------------------------|--------------------|------------------------|
| Locat | <u>iion</u> | Plt. Ht. inches | Test Wt. lbs/ac | Yield <u>lbs/ac</u> | Plt. Ht. <u>inches</u> | Test Wt. lbs/ac | Yield <u>lbs/ac</u> |
| 1990 | Bozeman,MT (irr) | 23 | 57 | 6431 | 32 | 53 | 4469 |
| | Conrad,MT #1 (dry) | 20 | 53 | 3052 | 27 | 50 | 2671 |
| 2 | Conrad,MT #2 (dry) | 19 | 55 | 2398 | 24 | 54 | 2236 |
| | Denton,MT (dry) | 20 | 49 | 2180 | 25 | 49 | 1744 |
| 1991 | Bozeman,MT (irr) | 33 | 55 | 5758 | 37 | 54 | 4174 |
| | Conrad,MT #1 (dry) | 27 | 55 | 2963 | 37 | 54 | 2498 |
| | Conrad,MT #2 (dry) | 23 | 57 | 4261 | 37 | 59 | 3893 |
| | Denton,MT (dry) | 19 | 58 | 2687 | 28 | 47 | 2183 |
| 1992 | Bozeman,MT (irr) | 24 | 55 | 4833 | 29 | 51 | 3905 |
| 1994 | Bozeman,MT (irr) | 24 | 54 | 6140 | 33 | 53 | 4810 |
| | Conrad,MT #1 (dry) | 21 | 59 | 3055 | 26 | 54 | 2680 |
| | Conrad,MT #2 (dry) | <u>22</u> | <u>58</u> | <u>3731</u> | <u>28</u> | <u>53</u> | <u>3538</u> |
| | mean | 22.9 | 55.4 | 3957 | 30.3 | 52.6 | 3233 |

الما الما الما

Table 3. Agronomic data comparing Merlin and WestBred Waxbar in USDA trials at Aberdeen, Idaho in 1994.

| | Heading Date <u>from Jan 1</u> | Plt.Ht inches | Test Wt. lbs/bu | Yield <u>bu/ac</u> |
|------------|--------------------------------------|------------------|--------------------|-----------------------|
| Merlin | 181 | 23 | 53.2 | 124 |
| W.B.Waxbar | 176 | 39 | 56.8 | 135 |

A25

Table 4. Heading date (from Jan. 1) of Merlin and check varieties in Montana State Univ. spring barley trials.

| Locations | | <u>M</u> | <u>ərlin</u> | <u>erlin</u> <u>Baronesse</u> | | <u>Gallatin</u> | |
|------------------|-----------|------------|--------------|-------------------------------|------------|-----------------|------------|
| | | <u>92</u> | 94 | <u>92</u> | 94 | 92 | 94 |
| Dryland | | | | | | | |
| · - | Havre | | 193 | | 191 | | 175 |
| * * | Sidney | 171 | 171 | 173 | 166 | 170 | 164 |
| | Moccasin | 174 | 179 | 171 | 176 | 170 | 174 |
| | Huntley | 162 | 170 | 162 | 168 | 159 | 167 |
| * | Conrad | | 182 | | 179 | | 173 |
| | Bozeman | 165 | 173 | 165 | 173 | 165 | 170 |
| Irrigated | | | | | | | |
| gatou | Kalispell | 164 | 178 | 160 | 175 | 158 | 176 |
| • | Sidney | 174 | 169 | 175 | 167 | 172 | 164 |
| | Huntley | 162 | 167 | 161 | 165 | 160 | 163 |
| - | Bozeman | <u>171</u> | <u>176</u> | <u>171</u> | <u>174</u> | <u>170</u> | <u>172</u> |
| | mean | 167.9 | 175.8 | 167.3 | 173.4 | 165.5 | 169.8 |

ng:

Table 5. Plant height in inches of Merlin and check varieties in Montana State Univ. spring barley trials.

| Locations | | Me | <u>erlin</u> | <u>Baro</u> | <u>nesse</u> | Gal | <u>llatin</u> |
|------------------|-----------|-------------|--------------|-------------|--------------|-------------|---------------|
| | | <u>92</u> | <u>94</u> | <u>92</u> | <u>94</u> | 92 | 94 |
| Dryland | | | | | | | |
| | Havre | | 18.2 | | 23.6 | | 28.6 |
| | Sidney | 21.5 | 15.8 | 27.5 | 23.0 | 28.8 | 27.2 |
| | Moccasin | 22.8 | 21.0 | 29.9 | 23.2 | 31.4 | 25.7 |
| | Huntley | 25.0 | 27.0 | 31.0 | 28.0 | 34.5 | 31.5 |
| | Conrad | | 22.0 | | 30.0 | | 35.0 |
| | Bozeman | 23.9 | 25.1 | 27.9 | 30.2 | 28.4 | 32.5 |
| Irrigated | | | | | | | |
| 3 | Conrad | | 25.0 | | 27.0 | | 34.0 |
| | Kalispell | 19.0 | 19.4 | 24.9 | 26.4 | 26.7 | 34.5 |
| | Sidney | 25.0 | 22.9 | 34.6 | 30.0 | 35.8 | 33.5 |
| 4. | Huntley | 26.0 | 28.0 | 35.5 | 28.0 | 39.5 | 31.5 |
| | Bozeman | <u>27.5</u> | <u>26.6</u> | 32.9 | 32.0 | <u>35.1</u> | 34.0 |
| | mean | 23.8 | 22.8 | 30.5 | 27.4 | 32.5 | 31.6 |

ne

Table 6. Test weight in pounds per bushel of Merlin and check varieties in Montana State Univ. spring barley trials.

| Locations | <u>Merlin</u> Bai | | <u>Baro</u> | <u>Baronesse</u> | | Gallatin | |
|------------------|-------------------|-------------|-------------|------------------|-------------|-------------|-------------|
| | | <u>92</u> | <u>94</u> | <u>92</u> | <u>94</u> | 92 | 94 |
| Dryland | | | | | | | |
| | Havre | | 42.5 | | 45.4 | | 46.6 |
| | Sidney | 60.7 | 59.6 | 55.5 | 53.5 | 54.3 | 53.0 |
| | Moccasin | 58.5 | 59.0 | 53.2 | 51.5 | 53.8 | 51.2 |
| | Huntley | 55.9 | 46.3 | 49.9 | 45.9 | 51.8 | 46.3 |
| | Conrad | | 50.7 | | 50.0 | | 52.4 |
| | Bozeman | 60.6 | 57.3 | 55.8 | 53.5 | 55.5 | 54.8 |
| Irrigated | | | | | | · | |
| 3 | Conrad | | 58.7 | | 54.7 | • | 54.7 |
| | Kalispell | 55.5 | 54.1 | 51.6 | 52.8 | 52.2 | 52.9 |
| | Sidney | 58.2 | 59.1 | 51.3 | 54.3 | 51.3 | 53.8 |
| | Huntley | 54.4 | 54.9 | 50.6 | 49.2 | 51.5 | 50.7 |
| | Bozeman | <u>60.8</u> | <u>59.4</u> | <u>56.4</u> | <u>54.1</u> | <u>55.5</u> | <u>55.4</u> |
| | mean | 58.1 | 54.7 | 53.0 | 51.4 | 53.2 | 52.0 |

NBL vi

Table 7. Yield in bushels per acre of Merlin and check varieties in Montana State Univ. spring barley trials.

| <u>Locations</u> | | <u>M</u> e | <u>erlin</u> | <u>Baronesse</u> | | <u>Gallatin</u> | |
|------------------|-----------|--------------|--------------|------------------|--------------|-----------------|--------------|
| | | <u>92</u> | <u>94</u> | <u>92</u> | <u>94</u> | <u>92</u> | <u>94</u> |
| Dryland | | | | | | | |
| | Havre | | 57.7 | | 68.8 | | 65.2 |
| | Sidney | 86.8 | 28.2 | 105.7 | 46.2 | 93.9 | 47.6 |
| | Moccasin | 73.2 | 46.6 | 87.9 | 53.0 | 83.4 | 47.9 |
| | Huntley | 113.2 | 45.3 | 123.9 | 57.3 | 125.0 | 54.9 |
| | Conrad | | 71.8 | | 92.3 | | 93.0 |
| | Bozeman | 114.3 | 104.0 | 133.7 | 121.8 | 109.0 | 109.9 |
| Irrigated | | | | | | | |
| ga.to a | Conrad | | 78.4 | | 92.9 | | 87.1 |
| | Kalispell | 85.9 | 62.5 | 123.8 | 92.0 | 108.3 | 96.5 |
| | Sidney | 125.4 | 69.0 | 151.6 | 99.6 | 116.2 | 101.3 |
| * | Huntley | 126.4 | 84.7 | 126.2 | 124.5 | 129.6 | 97.5 |
| | Bozeman | <u>141.6</u> | <u>138.3</u> | <u>160.3</u> | <u>137.7</u> | <u>137.6</u> | <u>123.4</u> |
| • | mean | 108.4 | 71.5 | 126.6 | 89.6 | 112.9 | 84.0 |

100

